arXiv:1208.1156v1 [nucl-th] 6 Aug 2012

Rotational properties of the superheavy nucleus ^{256}Rf in particle-number conserving cranked shell model

Zhen-Hua Zhang,[1,2] Jie Meng,[1,3,4] En-Guang Zhao,[1,2,5] and Shan-Gui Zhou[2,5,*]

[1] *State Key Laboratory of Nuclear Physics and Technology,*
School of Physics, Peking University, Beijing 100871, China
[2] *State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,*
Chinese Academy of Sciences, Beijing 100190, China
[3] *School of Physics and Nuclear Energy Engineering, Beihang University, Beijing 100191, China*
[4] *Department of Physics, University of Stellenbosch, Stellenbosch 7602, South Africa*
[5] *Center of Theoretical Nuclear Physics, National Laboratory of Heavy Ion Accelerator, Lanzhou 730000, China*

(Dated: August 7, 2012)

The kinematic and dynamic moments of inertia of the ground state band in the recently observed nucleus ^{256}Rf and some neighboring even-even nuclei are investigated by using a cranked shell model (CSM) with the pairing correlations treated by a particle-number conserving (PNC) method, in which the blocking effects are taken into account exactly. The determined spin of the observed ground state band in ^{256}Rf by comparing the kinematic moments of inertia with the calculations of the PNC-CSM agrees with previous spin assignment. The effects of the high order deformation ε_6 on the delayed alignment in ^{254}No relative to ^{256}Rf as well as on dynamic moment of inertia are discussed.

PACS numbers: 21.60.-n; 21.60.Cs; 23.20.Lv; 27.90.+b

In recent years, the in-beam spectroscopy of the nuclei with $Z \approx 100$ has become a hot topic [1–6]. These transfermium nuclei bring new information for the domain of superheavy nuclei. Experimental results show that these nuclei are well deformed. Due to deformation effects, the orbitals originating from the spherical subshells which are important to the next magic number may come close to the Fermi surface in these deformed nuclei. For example, the $\pi 1/2^-[521]$ and $\pi 3/2^-[521]$ orbitals are of particular interest since they stem from the spherical spin doublets $\pi 2f_{5/2,7/2}$ orbtials, whose splitting is very important to the location of the next proton shell closure. The high spin rotational states of these transfermium nuclei can give valuable information about the single particle orbitals near the Fermi surface, especially the high-j intruder orbitals ($\nu j_{15/2}$ or $\pi i_{13/2}$) which are sensitive to the Coriolis interaction.

In our previous work [7], the spectroscopy of the nuclei with $A \approx 250$ is systematically investigated by a particle-number conserving (PNC) method based on cranked shell model (CSM) [8, 9] with a new Nilsson parameter set which is obtained by fitting the experimental single-particle spectra in these nuclei. The calculated bandhead energies of the 1-quasiproton and 1-quasineutron bands in odd-A nuclei are improved dramatically comparing with those calculated by using the traditional Nilsson parameter [10]. The experimental kinematic moments of inertia (MOI's) for the rotational bands in even-even, odd-A, and odd-odd nuclei are reproduced quite well by the PNC-CSM calculations. In contrary to the conventional Bardeen-Cooper-Schrieffer (BCS) or Hartree-Fock-Bogolyubov (HFB) approach, in the PNC method, the CSM Hamiltonian is solved directly in a truncated Fock-space [11]. Therefore the particle-number is conserved and the Pauli blocking effects are taken into account exactly.

Quite recently, the ground state band (GSB) with spin up to $20\hbar$ was observed in the even-even superheavy nucleus ^{256}Rf ($Z = 104$) [12]. The two lowest-lying transitions of this rotational band are not observed in experiment due to the strong competition from internal conversion. Therefore the Harris 2-parameter expansion formula [13, 14] was used to extrapolate the energies of these transitions. It is worthwhile to mention that Rf is the first element whose stability is entirely derived from the shell correction energy, and this element marks the gateway to superheavy elements [15]. The spectrum and MOI's of ^{256}Rf can give information about the single-particle structure and the pairing interaction of the superheavy nuclei and provide a test for current nuclear models.

In this Rapid Communication, the PNC-CSM is used to investigate the rotational properties of ^{256}Rf in comparison with some neighboring even-even nuclei. The spin of the experimentally observed lowest-lying transition in the GSB is determined by comparing the MOI's extracted from different spin assignments with the calculations of the PNC-CSM.

The CSM Hamiltonian of an axially symmetric nucleus in the rotating frame can be written as

$$H_{\mathrm{CSM}} = H_0 + H_{\mathrm{P}} = H_{\mathrm{Nil}} - \omega J_x + H_{\mathrm{P}} \ , \qquad (1)$$

where H_{Nil} is the Nilsson Hamiltonian, $-\omega J_x$ is the Coriolis interaction with the cranking frequency ω about the x axis (perpendicular to the nuclear symmetry z axis).

---

[*] sgzhou@itp.ac.cn

$H_{\mathrm{P}} = H_{\mathrm{P}}(0) + H_{\mathrm{P}}(2)$ is the pairing interaction,

$$H_{\mathrm{P}}(0) = -G_0 \sum_{\xi\eta} a^\dagger_\xi a^\dagger_{\bar{\xi}} a_{\bar{\eta}} a_\eta \ , \tag{2}$$

$$H_{\mathrm{P}}(2) = -G_2 \sum_{\xi\eta} q_2(\xi) q_2(\eta) a^\dagger_\xi a^\dagger_{\bar{\xi}} a_{\bar{\eta}} a_\eta \ , \tag{3}$$

where $\bar{\xi}$ ($\bar{\eta}$) labels the time-reversed state of a Nilsson state ξ (η), $q_2(\xi) = \sqrt{16\pi/5}\langle\xi|r^2 Y_{20}|\xi\rangle$ is the diagonal element of the stretched quadrupole operator, and G_0 and G_2 are the effective strengths of monopole and quadrupole pairing interactions, respectively.

Instead of the usual single-particle level truncation in conventional shell-model calculations, a cranked many-particle configuration (CMPC) truncation (Fock space truncation) is adopted which is crucial to make the PNC calculations for low-lying excited states both workable and sufficiently accurate [16]. An eigenstate of H_{CSM} can be written as

$$|\Psi\rangle = \sum_i C_i \, |i\rangle \ , \qquad (C_i \text{ real}), \tag{4}$$

where $|i\rangle$ is a CMPC (an eigenstate of the one-body operator H_0). By diagonalizing the H_{CSM} in a sufficiently large CMPC space, sufficiently accurate solutions for low-lying excited eigenstates of H_{CSM} are obtained [7].

The kinematic and dynamic MOI for the state $|\Psi\rangle$ are

$$\begin{aligned} J^{(1)} &= \frac{1}{\omega} \langle\Psi|J_x|\Psi\rangle \\ &= \frac{1}{\omega}\left(\sum_i C_i^2 \, \langle i|J_x|i\rangle + 2\sum_{i<j} C_i C_j \, \langle i|J_x|j\rangle\right) \ , \end{aligned} \tag{5}$$

$$\begin{aligned} J^{(2)} &= \frac{\mathrm{d}}{\mathrm{d}\omega} \langle\Psi|J_x|\Psi\rangle \\ &= \frac{\mathrm{d}}{\mathrm{d}\omega}\left(\sum_i C_i^2 \, \langle i|J_x|i\rangle + 2\sum_{i<j} C_i C_j \, \langle i|J_x|j\rangle\right) . \end{aligned} \tag{6}$$

The parameters used in this work are all taken from Ref. [7]. The traditional Nilsson parameters (κ and μ) [10, 17] are optimized to reproduce the experimental level schemes for the rare-earth and actinide nuclei near the stability line. However, this parameter set can not describe well the experimental level schemes of transfermium nuclei. Therefore we proposed a new set of Nilsson parameters (κ and μ) in [7, 18] which was obtained by fitting the experimental single-particle spectra in these nuclei.

The experimental values of the deformation parameters for the transfermium nuclei are very scare and the predictions of different theories are not consistent with each other [19–21]. In the PNC-CSM calculations, the deformations are chosen to be close to existing experimental values and change smoothly with the proton and the neutron numbers. The deformation parameters of ^{256}Rf can be extrapolated from Table II in Ref. [7] as $\varepsilon_2 = 0.255$ and $\varepsilon_4 = 0.025$.

The CMPC space in this work is constructed in the proton $N = 4, 5, 6$ shells and the neutron $N = 6, 7$ shells. The dimensions of the CMPC space are about 1000 both for protons and neutrons. The effective pairing strengths are $G_p = 0.40$ MeV, $G_{2p} = 0.035$ MeV, $G_n = 0.30$ MeV, and $G_{2n} = 0.020$ MeV, which are the same for all even-even nuclei in this mass region (see Table III in Ref. [7]).

Figure 1 shows the calculated cranked Nilsson levels near the Fermi surface of ^{256}Rf. The positive (negative) parity levels are denoted by blue (red) lines. The signature $\alpha = +1/2$ ($\alpha = -1/2$) levels are denoted by solid (dotted) lines. From Fig. 1 it can be seen that there exist a proton gap at $Z = 100$ and a neutron gap at $N = 152$, which is consistent with the calculation by using Woods-Saxon potential [20, 22]. The $Z = 104$ proton energy spacing in our calculation is about 0.5 MeV, which is much larger than that of $Z = 102$. This situation is the opposite in Ref. [20]. Nevertheless, the $Z = 104$ gap in our calculation is not very significant comparing with that calculated by the self-consistent mean field models [21, 23], which is usually larger than 1 MeV. For protons, the sequence of single-particle levels near the Fermi surface in our calculation is quite similar with that determined from the experimental information of ^{255}Lr [24], in which the energies of $1/2^-[521]$ and $7/2^-[514]$ are nearly degenerate. For neutrons, it is shown experimentally that the ground state of ^{255}Rf is $9/2^-[734]$ [25], which is also consistent with our calculation. So the Nilsson levels used in our calculation are quite reasonable.

The kinematic MOI's depend sensitively on the spin assignment. In Fig. 2 we show the comparison of experimental kinematic MOI's of the GSB in ^{256}Rf extracted from different spin assignments with the calculations from PNC-CSM. The red up-triangles, black solid circles and blue down-triangles denote the experimental MOI's extracted by assigning the observed lowest-lying 161 keV transition as $8^+ \to 6^+$, $6^+ \to 4^+$, and $4^+ \to 2^+$, respectively. Our calculation agrees very well with the $6^+ \to 4^+$ assignment, and it is consistent with the spin assignment using the Harris formula in Ref. [12]. So in the following calculation, the 161 keV transition is assigned as $6^+ \to 4^+$ and the deduced energies of $4^+ \to 2^+$ (104 keV) and $2^+ \to 0^+$ (44 keV) in Ref. [12] are also used to extract the experimental kinematic and dynamic MOI's in the GSB of ^{256}Rf. It will be an interesting topic to make the spin assignment by using some other rotational formulae or models [26–33]. It is noted that spin assignment for the rotational bands in ^{253}No [34] have already made in Ref. [35] using a 2-parameter *ab* formula [26, 27].

In order to study the rotational properties of ^{256}Rf, the experimental (solid circles) and calculated (solid lines) kinematic MOI's $J^{(1)}$ for ^{256}Rf and the neighboring even-even nuclei ^{250}Fm [36] and ^{252,254}No [37, 38] are shown in Fig. 3. The results for the last three nuclei can also be found in our previous work [7]. It can be seen that the absolute values of the MOI's for these four nuclei are reproduced quite well, especially for ^{256}Rf. This again



FIG. 1. (Color online) The cranked Nilsson levels near the Fermi surface of ^{256}Rf (a) for protons and (b) for neutrons. The positive (negative) parity levels are denoted by blue (red) lines. The signature $\alpha = +1/2$ ($\alpha = -1/2$) levels are denoted by solid (dotted) lines.



FIG. 2. (Color online) The comparison of experimental kinematic MOI's of the GSB in ^{256}Rf extracted from different spin assignments with the PNC-CSM calculations. The red up-triangles, black solid circles and blue down-triangles denote the experimental MOI's extracted by assigning the 161 keV transition as from $8^+ \to 6^+$, $6^+ \to 4^+$, and $4^+ \to 2^+$, respectively. The data are taken from Ref. [12].



FIG. 3. The experimental (solid circles) and calculated (solid lines) kinematic MOI's $J^{(1)}$ for ^{256}Rf and the neighboring even-even nuclei ^{250}Fm [36] and ^{252,254}No [37, 38]. The calculated results for ^{250}Fm and ^{252,254}No can also be found in Ref. [7].

indicates that the single-particle levels we adopted are reasonable in this mass region, which shows that there exist a proton gap at $Z = 100$ and a neutron gap at $N = 152$ and there is no significant proton gap at $Z = 104$.

In Fig. 4 we show the experimental (solid circles) and calculated (solid black lines) dynamic MOI's $J^{(2)}$ for ^{256}Rf and the neighboring even-even nuclei ^{250}Fm [36] and ^{252,254}No [37, 38]. The experimental dynamic MOI's $J^{(2)}$ for ^{256}Rf are reproduced perfectly by the PNC-CSM calculation. For the other nuclei the results are quite reasonable although some deviations exist. As pointed out in Ref. [12], the alignment of $N = 150$ isotones (^{250}Fm and ^{252}No) occurs a little earlier than that of $N = 152$ isotones (^{254}No and ^{256}Rf) and is delayed in ^{254}No relative to ^{256}Rf. The upbending mechanism in this mass region has been investigated in detail in our previous work [7]. Similar results have been achieved by other models [21, 39]. However, we can not reproduce the alignment delay in ^{254}No. In Ref. [40], this phenomenon is explained in terms of β_6 deformation which decreases the energies of the neutron $j_{15/2}$ intruder orbitals below



FIG. 4. (Color online) The experimental (solid circles) and calculated (solid black lines) dynamic MOI's $J^{(2)}$ for ^{256}Rf and the neighboring even-even nuclei ^{250}Fm [36] and ^{252,254}No [37, 38]. The red lines are the results when the high order deformation parameter ε_6 is considered in the PNC-CSM calculation. ε_6 for ^{250}Fm, ^{252,254}No, and ^{256}Rf are 0.044, 0.040, 0.042, and 0.038, respectively, which are taken from Ref. [19].

the $N = 152$ gap. Here in Fig. 4 we show our results after considering this high order deformation. The red lines are the results when ε_6 is considered in the PNC-CSM calculation. ε_6 for ^{250}Fm, ^{252,254}No, and ^{256}Rf are 0.044, 0.040, 0.042, and 0.038, respectively, which are taken from Ref. [19]. It can be seen that the ε_6 deformation have prominent effect in the higher spin region. The results are improved after considering this deformation in the PNC-CSM. Note that the deformation parameter ε_6 is fixed during our calculation while it is changing with the rotational frequency in the Total Routhian Surface (TRS) calculation in Ref. [40]. We expect that after considering this effect, the results can be improved further.

In summary, the recently observed high-spin rotational GSB in ^{256}Rf is analyzed by using a cranked shell model with the pairing correlations treated by a particle-number conserving method in which the blocking effects are taken into account exactly. Both the experimental kinematic and dynamic MOI's are reproduced quite well by the PNC-CSM calculations. The spin of the experimentally observed lowest-lying transition in the GSB is determined by comparing the MOI's extracted from different spin assignments with the calculations of the PNC-CSM. The rotational property differences between ^{256}Rf and some neighboring even-even nuclei are illustrated. The effects of the high order deformation ε_6 on the delayed alignment in ^{254}No relative to ^{256}Rf is discussed.

We thank P. T. Greenlees for sending us the data of ^{256}Rf prior to publication. Helpful discussions with P. T. Greenlees, J. Piot, and P. W. Zhao are gratefully acknowledged. This work has been supported by the Major State Basic Research Development Program of China (973 Program: "New physics and technology at the limits of nuclear stability"), National Natural Science Foundation of China (Grant No. 10975008, No. 10975100, No. 10979066, No. 11121403, No. 11175002, No. 11175252, and No. 11120101005), Knowledge Innovation Project of Chinese Academy of Sciences (Grant No. KJCX2-EW-N01 and No. KJCX2-YW-N32), the Research Fund for the Doctoral Program of Higher Education under Grant No. 20110001110087. The computation of this work was supported by Supercomputing Center, CNIC of CAS. The results described in this paper are obtained on the ScGrid of Supercomputing Center, Computer Network Information Center of Chinese Academy of Sciences.

---

[1] M. Leino and F. Heßberger, Annu. Rev. Nucl. Part. Sci. **54**, 175 (2004).
[2] D. Ackermann, Acta Phys. Pol. B **42**, 577 (2011).
[3] R.-D. Herzberg, J. Phys. G: Nucl. Part. Phys. **30**, R123 (2004).
[4] R.-D. Herzberg and P. Greenlees, Prog. Part. Nucl. Phys. **61**, 674 (2008).
[5] P. T. Greenlees, Acta Phys. Pol. B **42**, 587 (2011).
[6] J. Piot, B. J.-P. Gall, O. Dorvaux, P. T. Greenlees, N. Rowley, L. L. Andersson, D. M. Cox, F. Dechery, T. Grahn, K. Hauschild, G. Henning, A. Herzan, R.-D. Herzberg, F. P. Heßberger, U. Jakobsson, P. Jones, R. Julin, S. Juutinen, S. Ketelhut, T.-L. Khoo, M. Leino, J. Ljungvall, A. Lopez-Martens, P. Nieminen, J. Pakarinen, P. Papadakis, E. Parr, P. Peura, P. Rahkila, S. Rinta-Antila, J. Rubert, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, D. Seweryniak, J. Sorri, B. Sulignano, and J. Uusitalo, Phys. Rev. C **85**, 041301R (2012).
[7] Z.-H. Zhang, X.-T. He, J.-Y. Zeng, E.-G. Zhao, and S.-G. Zhou, Phys. Rev. C **85**, 014324 (2012).
[8] J. Y. Zeng and T. S. Cheng, Nucl. Phys. A **405**, 1 (1983).
[9] J. Y. Zeng, T. H. Jin, and Z. J. Zhao, Phys. Rev. C **50**, 1388 (1994).
[10] S. G. Nilsson, C. F. Tsang, A. Sobiczewski, Z. Szymaski, S. Wycech, C. Gustafson, I. L. Lamm, P. Möller, and B. Nilsson, Nucl. Phys. A **131**, 1 (1969).
[11] C. S. Wu and J. Y. Zeng, Phys. Rev. C **39**, 666 (1989).
[12] P. T. Greenlees, J. Rubert, J. Piot, B. J. P. Gall, L. L. Andersson, M. Asai, Z. Asfari, D. M. Cox, F. Dechery, O. Dorvaux, T. Grahn, K. Hauschild, G. Henning, A. Herzan, R.-D. Herzberg, F. P. Heßberger, U. Jakobsson, P. Jones, R. Julin, S. Juutinen, S. Ketelhut, T.-L. Khoo, M. Leino, J. Ljungvall, A. Lopez-Martens, R. Lozeva, P. Nieminen, J. Pakarinen, P. Papadakis, E. Parr, P. Peura, P. Rahkila, S. Rinta-Antila, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, D. Seweryniak, J. Sorri, B. Sulignano, C. Theisen, J. Uusitalo, and M. Venhart, Phys. Rev. Lett. **109**, 012501 (2012).
[13] S. M. Harris, Phys. Rev. Lett. **13**, 663 (1964).

[14] S. M. Harris, Phys. Rev. **138**, B509 (1965).

[15] M. Schädel, Angew. Chem. Int. Ed. **45**, 368 (2006).

[16] H. Molique and J. Dudek, Phys. Rev. C **56**, 1795 (1997).

[17] T. Bengtsson and I. Ragnarsson, Nucl. Phys. A **436**, 14 (1985).

[18] Z.-H. Zhang, J.-Y. Zeng, E.-G. Zhao, and S.-G. Zhou, Phys. Rev. C **83**, 011304R (2011).

[19] P. Möller and J. Nix, At. Data and Nucl. Data Tables **59**, 185 (1995).

[20] A. Sobiczewski, I. Muntian, and Z. Patyk, Phys. Rev. C **63**, 034306 (2001).

[21] A. V. Afanasjev, T. L. Khoo, S. Frauendorf, G. A. Lalazissis, and I. Ahmad, Phys. Rev. C **67**, 024309 (2003).

[22] R. R. Chasman, I. Ahmad, A. M. Friedman, and J. R. Erskine, Rev. Mod. Phys. **49**, 833 (1977); Rev. Mod. Phys. **50**, 173 (1978).

[23] M. Bender, P. Bonche, T. Duguet, and P. H. Heenen, Nucl. Phys. A **723**, 354 (2003).

[24] A. Chatillon, C. Theisen, P. Greenlees, G. Auger, J. Bastin, E. Bouchez, B. Bouriquet, J. Casandjian, R. Cee, E. Clément, R. Dayras, G. de France, R. de Toureil, S. Eeckhaudt, A. Görgen, T. Grahn, S. Grévy, K. Hauschild, R. Herzberg, P. Ikin, G. Jones, P. Jones, R. Julin, S. Juutinen, H. Kettunen, A. Korichi, W. Korten, Y. Le Coz, M. Leino, A. Lopez-Martens, S. Lukyanov, Y. Penionzhkevich, J. Perkowski, A. Pritchard, P. Rahkila, M. Rejmund, J. Saren, C. Scholey, S. Siem, M. Saint-Laurent, C. Simenel, Y. Sobolev, C. Stodel, J. Uusitalo, A. Villari, M. Bender, P. Bonche, and P. Heenen, Eur. Phys. J. A **30**, 397 (2006).

[25] F. Heßberger, S. Hofmann, D. Ackermann, S. Antalic, B. Kindler, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, K. Nishio, A. Popeko, B. Sulignano, S. Saro, B. Streicher, M. Venhart, and A. Yeremin, Eur. Phys. J. A **30**, 561 (2006).

[26] P. Holmberg and P. Lipas, Nucl. Phys. A **117**, 552 (1968).

[27] C. S. Wu and J. Y. Zeng, Commu. Theor. Phys. **8**, 51 (1987).

[28] J. Y. Zeng, J. Meng, C. S. Wu, E. G. Zhao, Z. Xing, and X. Q. Chen, Phys. Rev. C **44**, 1745R (1991).

[29] C. S. Wu, J. Y. Zeng, Z. Xing, X. Q. Chen, and J. Meng, Phys. Rev. C **45**, 261 (1992).

[30] F. Xu and J. Hu, Phys. Rev. C **49**, 1449 (1994).

[31] F. Xu and J. Hu, Phys. Rev. C **52**, 431 (1995).

[32] S.-G. Zhou, C. Zheng, F. Xu, and J. Hu, Nucl. Phys. A **615**, 229 (1997).

[33] S.-G. Zhou and C. Zheng, Phys. Rev. C **55**, 2324 (1997).

[34] R. Herzberg, S. Moon, S. Eeckhaudt, P. Greenlees, P. Butler, T. Page, A. Afanasjev, N. Amzal, J. Bastin, F. Becker, M. Bender, B. Bruyneel, J. Cocks, I. Darby, O. Dorvaux, K. Eskola, J. Gerl, T. Grahn, C. Gray-Jones, N. Hammond, K. Hauschild, P. Heenen, K. Helariutta, A. Herzberg, F. Hessberger, M. Houry, A. Hürstel, R. Humphreys, G. Jones, P. Jones, R. Julin, S. Juutinen, H. Kankaanpȧȧ, H. Kettunen, T. Khoo, W. Korten, P. Kuusiniemi, Y. LeCoz, M. Leino, A. Leppȧnen, C. Lister, R. Lucas, M. Muikku, P. Nieminen, M. Nyman, R. Page, T. Page, J. Pakarinen, A. Pritchard, P. Rahkila, P. Reiter, M. Sandzelius, J. Saren, C. Schlegel, C. Scholey, C. Theisen, W. Trzaska, J. Uusitalo, A. Wiens, and H. Wollersheim, Eur. Phys. J. A **42**, 333 (2009).

[35] K. Wen, Z.-H. Zhang, E.-G. Zhao, and S.-G. Zhou, Sci. Sin.-Phys. Mech. Astron. **42**, 22 (2012), in Chinese.

[36] J. E. Bastin, R.-D. Herzberg, P. A. Butler, G. D. Jones, R. D. Page, D. G. Jenkins, N. Amzal, P. M. T. Brew, N. J. Hammond, R. D. Humphreys, P. J. C. Ikin, T. Page, P. T. Greenlees, P. M. Jones, R. Julin, S. Juutinen, H. Kankaanpää, A. Keenan, H. Kettunen, P. Kuusiniemi, M. Leino, A. P. Leppänen, M. Muikku, P. Nieminen, P. Rahkila, C. Scholey, J. Uusitalo, E. Bouchez, A. Chatillon, A. Hürstel, W. Korten, Y. L. Coz, C. Theisen, D. Ackermann, J. Gerl, K. Helariutta, F. P. Hessberger, C. Schlegel, H. J. Wollerscheim, M. Lach, A. Maj, W. Meczynski, J. Styczen, T. L. Khoo, C. J. Lister, A. V. Afanasjev, H. J. Maier, P. Reiter, P. Bednarczyc, K. Eskola, and K. Hauschild, Phys. Rev. C **73**, 024308 (2006).

[37] R.-D. Herzberg, N. Amzal, F. Becker, P. A. Butler, A. J. C. Chewter, J. F. C. Cocks, O. Dorvaux, K. Eskola, J. Gerl, P. T. Greenlees, N. J. Hammond, K. Hauschild, K. Helariutta, F. Heßberger, M. Houry, G. D. Jones, P. M. Jones, R. Julin, S. Juutinen, H. Kankaanpää, H. Kettunen, T. L. Khoo, W. Korten, P. Kuusiniemi, Y. L. Coz, M. Leino, C. J. Lister, R. Lucas, M. Muikku, P. Nieminen, R. D. Page, P. Rahkila, P. Reiter, C. Schlegel, C. Scholey, O. Stezowski, C. Theisen, W. H. Trzaska, J. Uusitalo, and H. J. Wollersheim, Phys. Rev. C **65**, 014303 (2001).

[38] S. Eeckhaudt, P. Greenlees, N. Amzal, J. Bastin, E. Bouchez, P. Butler, A. Chatillon, K. Eskola, J. Gerl, T. Grahn, A. Görgen, R. Herzberg, F. Hessberger, A. Hürstel, P. Ikin, G. Jones, P. Jones, R. Julin, S. Juutinen, H. Kettunen, T. Khoo, W. Korten, P. Kuusiniemi, Y. Le Coz, M. Leino, A. Leppänen, P. Nieminen, J. Pakarinen, J. Perkowski, A. Pritchard, P. Reiter, P. Rahkila, C. Scholey, C. Theisen, J. Uusitalo, K. Van de Vel, and J. Wilson, Eur. Phys. J. A **26**, 227 (2005).

[39] F. Al-Khudair, G.-L. Long, and Y. Sun, Phys. Rev. C **79**, 034320 (2009).

[40] H. L. Liu, F. R. Xu, and P. M. Walker, Phys. Rev. C **86**, 011301R (2012).